arXiv:1409.5959v3 [math.CO] 9 May 2015

Automorphism group of the modified bubble-sort graph

Ashwin Ganesan*

Abstract

The modified bubble-sort graph of dimension n is the Cayley graph of S_n generated by n cyclically adjacent transpositions. In the present paper, it is shown that the automorphism group of the modified bubble sort graph of dimension n is $S_n \times D_{2n}$, for all $n \geq 5$. Thus, a complete structural description of the automorphism group of the modified bubble-sort graph is obtained. A similar direct product decomposition is seen to hold for arbitrary normal Cayley graphs generated by transposition sets.

Index terms — modified bubble-sort graph; automorphism group; Cayley graphs; transposition sets.

1. Introduction

Let $X = (V, E)$ be a simple undirected graph. The (full) automorphism group of X, denoted by $\mathrm{Aut}(X)$, is the set of permutations of the vertex set that preserves adjacency, i.e., $\mathrm{Aut}(X) := \{g \in \mathrm{Sym}(V) : E^g = E\}$. Let H be a group with identity element e, and let S be a subset of H. The Cayley graph of H with respect to S, denoted by $\mathrm{Cay}(H, S)$, is the graph with vertex set H and arc set $\{(h, sh) : h \in H, s \in S\}$. When S satisfies the condition $1 \notin S = S^{-1}$, the Cayley graph $\mathrm{Cay}(H, S)$ has no self-loops and can be considered to be undirected.

A Cayley graph $\mathrm{Cay}(H, S)$ is vertex-transitive since the right regular representation $R(H)$ acts as a group of automorphisms of the Cayley graph. The set of automorphisms of H that fixes S setwise is a subgroup of the stabilizer $\mathrm{Aut}(\mathrm{Cay}(H, S))_e$ (cf. [1], [7]). A Cayley graph $X := \mathrm{Cay}(H, S)$ is said to be *normal* if $R(H)$ is a normal subgroup of $\mathrm{Aut}(X)$, or equivalently, if $\mathrm{Aut}(X) = R(H) \rtimes \mathrm{Aut}(H, S)$ (cf. [9]).

Let S be a set of transpositions generating the symmetric group S_n. The transposition graph of S, denoted by $T(S)$, is defined to be the graph with vertex set $\{1, \ldots, n\}$, and with two vertices i and j being adjacent in $T(S)$ whenever $(i, j) \in S$. A set S of

*Department of Electronics and Telecommunication Engineering, Vidyalankar Institute of Technology, Wadala, Mumbai, India. Correspondence address: ashwin.ganesan@gmail.com

transpositions generates S_n iff the transposition graph of S is connected. When the transposition graph of S is the n-cycle graph, then the Cayley graph $\mathrm{Cay}(S_n, S)$ is called the modified bubble-sort graph of dimension n. Thus, the modified bubble-sort graph of dimension n is the Cayley graph of S_n with respect to the set of generators $\{(1,2),(2,3),\ldots,(n-1,n),(n,1)\}$. The modified bubble-sort graph has been investigated for consideration as the topology of interconnection networks (cf. [8]). Many authors have investigated the automorphism group of graphs that arise as the topology of interconnection networks; for example, see [2], [3], [5], [10], [11].

Godsil and Royle [7] showed that if the transposition graph of S is an asymmetric tree, then the automorphism group of the Cayley graph $\mathrm{Cay}(S_n, S)$ is isomorphic to S_n. Feng [4] showed that $\mathrm{Aut}(S_n, S)$ is isomorphic to $\mathrm{Aut}(T(S))$ and that if the transposition graph of S is an arbitrary tree, then the automorphism group of $\mathrm{Cay}(S_n, S)$ is the semidirect product $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$. Ganesan [6] showed that if the girth of the transposition graph of S is at least 5, then the automorphism group of the Cayley graph $\mathrm{Cay}(S_n, S)$ is the semidirect product $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$. The results in the present paper imply that all these automorphism groups in the literature can be factored as a direct product.

In Zhang and Huang [10], it was shown the automorphism group of the modified bubble-sort graph of dimension n is the group product $S_n D_{2n}$ (groups products are also referred to as Zappa-Szep products). This result was strengthened in Feng [4], where it was proved that the automorphism group of the modified bubble-sort graph of dimension n is the semidirect product $R(S_n) \rtimes D_{2n}$ (cf. [4, p. 72] for an explicit statement of this conclusion).

In the present paper, we obtain a complete structural description of the automorphism group of the modified bubble-sort graph of dimension n:

Theorem 1. *The automorphism group of the modified bubble-sort graph of dimension n is $S_n \times D_{2n}$, for all $n \geq 5$.*

We shall prove the following more general result:

Theorem 2. *Let S be a set of transpositions generating $S_n (n \geq 3)$ such that the Cayley graph $\mathrm{Cay}(S_n, S)$ is normal. Then, the automorphism group of the Cayley graph $\mathrm{Cay}(S_n, S)$ is the direct product $S_n \times \mathrm{Aut}(T(S))$, where $T(S)$ denotes the transposition graph of S.*

In the special case where $T(S)$ is the n-cycle graph, $\mathrm{Aut}(T(S))$ is isomorphic to the dihedral group D_{2n} of order $2n$. Hence, Theorem 1 is a special case of Theorem 2. Also, Ganesan [6] showed that the modified bubble-sort graphs of dimension less than 5 are non-normal; hence, the assumption $n \geq 5$ in Theorem 1 is necessary.

Remark 1. Given a set S of transpositions generating S_n, let $G := \mathrm{Aut}(\mathrm{Cay}(S_n, S))$. In the instances where $G = R(S_n) \rtimes G_e$, the factor $G_e \cong \mathrm{Aut}(T(S))$ is in general not a normal subgroup of G, and so the semidirect product cannot be written immediately as a direct product. For example, for the modified bubble-sort graph of dimension n, $G \cong R(S_n) \rtimes G_e \cong S_n \rtimes D_{2n}$, where G_e is not normal in G. In the present paper, it is

shown that $R(S_n)$ has another complement in G which is a normal subgroup of G. In the proof below, we show that the image of $\mathrm{Aut}(T(S))$ under the left regular action of S_n on itself is a normal complement of $R(S_n)$ in G. Thus, the direct factor $\mathrm{Aut}(T(S))$ that arises in $G \cong R(S_n) \times \mathrm{Aut}(T(S))$ is not G_e but is obtained in a different manner.

2. Proof of Theorem 2

Let S be a set of transpositions generating S_n. We first establish that the Cayley graph $\mathrm{Cay}(S_n, S)$ has a particular subgroup of automorphisms. In this section, let λ denote the left regular action of S_n on itself, defined by $\lambda : S_n \to \mathrm{Sym}(S_n), a \mapsto \lambda_a$, where $\lambda_a : x \mapsto a^{-1}x$.

Proposition 3. *Let $T(S)$ denote the transposition graph of S. Then, $\{\lambda_a : a \in \mathrm{Aut}(T(S))\}$ is a set of automorphisms of the Cayley graph $X := \mathrm{Cay}(S_n, S)$.*

Proof: Let $a \in \mathrm{Aut}(T(S))$. We show that $\{h, g\} \in E(X)$ if and only if $\{h, g\}^{\lambda(a)} \in E(X)$. Suppose $\{h, g\} \in E(X)$. Then $g = sh$ for some transposition $s = (i, j) \in S$. We have that $\{h, g\}^{\lambda(a)} = \{h, sh\}^{\lambda(a)} = \{h^{\lambda(a)}, (sh)^{\lambda(a)}\} = \{a^{-1}h, a^{-1}sh\} = \{a^{-1}h, (a^{-1}sa)a^{-1}h\}$. Now $a^{-1}sa = a^{-1}(i, j)a = (i^a, j^a) \in S$ since a is an automorphism of the graph $T(S)$ that has edge set S. Thus, $\{h, sh\}^{\lambda(a)} \in E(X)$. Conversely, suppose $\{h, g\}^{\lambda(a)} \in E(X)$. Then $a^{-1}h = sa^{-1}g$ for some $s \in S$. Hence $h = (asa^{-1})g$. We have that $asa^{-1} = a(i, j)a^{-1} = (i, j)^{a^{-1}} \in S$ because a is an automorphism of $T(S)$. Hence, h is adjacent to g. Thus, $\lambda(\mathrm{Aut}(T(S)))$ is a subgroup of $\mathrm{Aut}(X)$. ∎

Theorem 4. *Let S be a set of transpositions generating $S_n (n \geq 3)$ such that the Cayley graph $\mathrm{Cay}(S_n, S)$ is normal. Then, the automorphism group of the Cayley graph $\mathrm{Cay}(S_n, S)$ is $S_n \times \mathrm{Aut}(T(S))$, where $T(S)$ denotes the transposition graph of S.*

Proof: Let X denote the Cayley graph $\mathrm{Cay}(S_n, S)$. Since X is a normal Cayley graph, its automorphpism group $\mathrm{Aut}(X)$ is equal to $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$ (cf. [9]). Let $R(a)$ denote the permutation of S_n induced by action by right multiplication by a, so that $R(S_n) := \{R(a) : a \in S_n\}$ is the right regular representation of S_n. The intersection of the left and right regular representations of a group is the image of the center of the group under either action. The center of S_n is trivial, whence $R(S_n) \cap \lambda(S_n) = 1$. In particular, $\lambda(\mathrm{Aut}(T(S)))$ and $R(S_n)$ have a trivial intersection. By Feng [4], $\mathrm{Aut}(S_n, S) \cong \mathrm{Aut}(T(S))$, and it follows from cardinality arguments that $R(S_n)\lambda(\mathrm{Aut}(T(S)))$ exhausts all the elements of $\mathrm{Aut}(X)$. Thus, $R(S_n)$ and $\lambda(\mathrm{Aut}(T(S)))$ are complements of each other in $\mathrm{Aut}(X)$ and every element in $\mathrm{Aut}(X)$ can be expressed uniquely in the form $R(a)\lambda(b)$ for some $a \in S_n$ and $b \in \mathrm{Aut}(T(S))$. This proves that $\mathrm{Aut}(X) = R(S_n) \rtimes \lambda(\mathrm{Aut}(T(S)))$.

It remains to prove that $\lambda(\mathrm{Aut}(T(S)))$ is a normal subgroup of $\mathrm{Aut}(X)$. Suppose $g \in \mathrm{Aut}(X)$ and $c \in \mathrm{Aut}(T(S))$. We show that $g^{-1}\lambda(c)g \in \lambda(\mathrm{Aut}(T(S)))$. We have that $g = R(a)\lambda(b)$ for some $a \in S_n, b \in \mathrm{Aut}(T(S))$. Hence, $g^{-1}\lambda(c)g = (R(a)\lambda(b))^{-1}\lambda(c)(R(a)\lambda(b))$, which maps $x \in S_n$ to $b^{-1}c^{-1}bxa^{-1}a = b^{-1}c^{-1}bx$. Since $b, c \in \mathrm{Aut}(T(S)), d^{-1} := b^{-1}c^{-1}b \in \mathrm{Aut}(T(S))$. Thus, $g^{-1}\lambda(c)g = \lambda(d) \in \lambda(\mathrm{Aut}(T(S)))$.

Hence, $\lambda(\mathrm{Aut}(T(S)))$ is a normal subgroup of $\mathrm{Aut}(X)$ and $\mathrm{Aut}(X) = R(S_n) \times \lambda(\mathrm{Aut}(T(S)))$. Since $\lambda(\mathrm{Aut}(T(S))) \cong \mathrm{Aut}(T(S))$, the assertion follows. ∎

Remark 2. We recall a particular result from group theory, which can be used to deduce that the semidirect products in the literature can be strengthened to direct products. Let A be a subgroup of a group H and suppose H has a trivial center. Let A act on H by conjugation. Let $\lambda(A)$ denote the image of the left action of A on H. Then the groups $R(H) \rtimes \mathrm{Inn}(A)$ and $R(H) \times \lambda(A)$ are isomorphic, where both groups are internal group products and subgroups of $\mathrm{Sym}(H)$. It follows from this group-theoretic result that the automorphism group of the Cayley graphs mentioned above can be factored as direct products. However, to the best of our knowledge, this group-theoretic result has not been used so far to deduce results in the context of automorphism groups of Cayley graphs generated by transposition sets - the expressions given in the previous literature for the automorphism group of Cayley graphs mentioned above have been only semidirect product factorizations (cf. [4, p.72], [6], [9]). In the present paper, in addition to obtaining a complete structural description of the automorphism group of the modified bubble-sort graph and of a family of normal Cayley graphs, the proof method also includes Proposition 3, which establishes that these graphs possess certain automorphisms.

References

[1] N. L. Biggs. *Algebraic Graph Theory, 2nd Edition*. Cambridge University Press, Cambridge, 1993.

[2] Y.-P. Deng and X.-D. Zhang. Automorphism group of the derangement graph. *The Electronic Journal of Combinatorics*, 18:#P198, 2011.

[3] Y.-P. Deng and X.-D. Zhang. Automorphism groups of the pancake graphs. *Information Processing Letters*, 112:264–266, 2012.

[4] Y-Q. Feng. Automorphism groups of Cayley graphs on symmetric groups with generating transposition sets. *Journal of Combinatorial Theory Series B*, 96:67–72, 2006.

[5] A. Ganesan. Automorphism group of the complete transposition graph. *Journal of Algebraic Combinatorics, to appear*.

[6] A. Ganesan. Automorphism groups of Cayley graphs generated by connected transposition sets. *Discrete Mathematics*, 313:2482–2485, 2013.

[7] C. Godsil and G. Royle. *Algebraic Graph Theory*. Graduate Texts in Mathematics vol. 207, Springer, New York, 2001.

[8] S. Lakshmivarahan, J-S. Jho, and S. K. Dhall. Symmetry in interconnection networks based on Cayley graphs of permutation groups: A survey. *Parallel Computing*, 19:361–407, 1993.

[9] M. Y. Xu. Automorphism groups and isomorphisms of Cayley digraphs. *Discrete Mathematics*, 182:309–319, 1998.

[10] Z. Zhang and Q. Huang. Automorphism groups of bubble sort graphs and modified bubble sort graphs. *Advances in Mathematics (China)*, 34(4):441–447, 2005.

[11] J-X. Zhou. The automorphism group of the alternating group graph. *Applied Mathematics Letters*, 24:229–231, 2011.